

# The Arbelos in $n$ -Aliquot Parts

Hiroshi Okumura and Masayuki Watanabe

**Abstract.** We generalize the classical arbelos to the case divided into many chambers by semicircles and construct embedded patterns of such arbelos.

## 1. Introduction and preliminaries

Let  $\{\alpha, \beta, \gamma\}$  be an arbelos, that is,  $\alpha, \beta, \gamma$  are semicircles whose centers are collinear and erected on the same side of this line,  $\alpha, \beta$  are tangent externally, and  $\gamma$  touches  $\alpha$  and  $\beta$  internally. In this paper we generalize results on the Archimedean circles of the arbelos. We take the line passing through the centers of  $\alpha, \beta, \gamma$  as the  $x$ -axis and the line passing through the tangent point  $O$  of  $\alpha$  and  $\beta$  and perpendicular to the  $x$ -axis as the  $y$ -axis. Let  $\alpha_0 = \alpha, \alpha_1, \dots, \alpha_n = \beta$  be  $n + 1$  distinct semicircles touching  $\alpha$  and  $\beta$  at  $O$ , where  $\alpha_1, \dots, \alpha_{n-1}$  are erected on the same side as  $\alpha$  and  $\beta$ , and intersect with  $\gamma$ . One of them may be the line perpendicular to the  $x$ -axis (i.e.  $y$ -axis). If the  $n$  inscribed circles in the curvilinear triangles bounded by  $\alpha_{i-1}, \alpha_i, \gamma$  are congruent we call this configuration of semicircles  $\{\alpha_0 = \alpha, \alpha_1, \dots, \alpha_n = \beta, \gamma\}$  an arbelos in  $n$ -aliquot parts, and the inscribed circles the Archimedean circles in  $n$ -aliquot parts. In this paper we calculate the radii of the Archimedean circles in  $n$ -aliquot parts and construct embedded patterns of arbelos in aliquot parts.




Figure 1. The case  $n = 3$

For the arbelos  $\{\alpha, \beta, \gamma\}$  we denote by  $\Phi(\alpha, \beta, \gamma)$  the family of semicircles through  $O$ , having the common point with  $\gamma$  in the region  $y \geq 0$  and with centers on the  $x$ -axis, together with the line perpendicular to the  $x$ -axis at  $O$ . Renaming if necessary we assume  $\alpha$  in the region  $x \geq 0$ . Let  $a, b$  be the radii of  $\alpha, \beta$ . The semicircle  $\gamma$  meets the  $x$ -axis at  $-2b$  and  $2a$ .

For a semicircle  $\alpha_i \in \Phi(\alpha, \beta, \gamma)$ , let  $a_i$  be the  $x$ -coordinate of its center. Define  $\mu(\alpha_i)$  as follows.

If  $a \neq b$ ,

$$\mu(\alpha_i) = \begin{cases} \frac{a_i - a + b}{a_i}, & \text{if } \alpha_i \text{ is a semi-circle,} \\ 1, & \text{if } \alpha_i \text{ is the line.} \end{cases}$$

If  $a = b$ ,

$$\mu(\alpha_i) = \begin{cases} \frac{1}{a_i}, & \text{if } \alpha_i \text{ is a semi-circle,} \\ 0, & \text{if } \alpha_i \text{ is the line.} \end{cases}$$

In both cases  $\mu(\alpha_i)$  depends only on  $\alpha_i$  and the center of  $\gamma$ , but not on the radius of  $\gamma$ . For  $\alpha_i, \alpha_j \in \Phi(\alpha, \beta, \gamma)$ , the equality  $\mu(\alpha_i) = \mu(\alpha_j)$  holds if and only if  $\alpha_i = \alpha_j$ . For any  $\alpha_i \in \Phi(\alpha, \beta, \gamma)$ ,

$$\begin{aligned} \frac{b}{a} = \mu(\alpha) \geq \mu(\alpha_i) \geq \mu(\beta) = \frac{a}{b} & \text{ if } a < b, \\ \frac{1}{a} = \mu(\alpha) \geq \mu(\alpha_i) \geq \mu(\beta) = -\frac{1}{a} & \text{ if } a = b, \\ \frac{b}{a} = \mu(\alpha) \leq \mu(\alpha_i) \leq \mu(\beta) = \frac{a}{b} & \text{ if } a > b. \end{aligned}$$

For  $\alpha_i, \alpha_j \in \Phi(\alpha, \beta, \gamma)$ , define the order

$$\alpha_i < \alpha_j \text{ if and only if } \begin{cases} \mu(\alpha_i) > \mu(\alpha_j) & \text{if } a \leq b, \\ \mu(\alpha_i) < \mu(\alpha_j) & \text{otherwise.} \end{cases}$$

This means that  $\alpha_i$  is nearer to  $\alpha$  than  $\alpha_j$  is. Throughout this paper we shall adopt these notations and assumptions.

## 2. An arbelos in aliquot parts

**Lemma 1.** *If  $\alpha_i$  and  $\alpha_j$  are semicircles in  $\Phi(\alpha, \beta, \gamma)$  with  $\alpha_i < \alpha_j$ , the radius of the inscribed circle in the curvilinear triangle bounded by  $\alpha_i, \alpha_j$  and  $\gamma$  is*

$$\frac{ab(a_j - a_i)}{a_i a_j - a a_i + b a_j}.$$

*Proof.* Let  $\mathcal{C}$  be the inscribed circle with radius  $r$ . First we invert  $\{\alpha_i, \alpha_j, \gamma, \mathcal{C}\}$  in the circle with center  $O$  and radius  $k$ . Then  $\alpha_i$  and  $\alpha_j$  are inverted to the lines  $\overline{\alpha_i}$  and  $\overline{\alpha_j}$  perpendicular to the  $x$ -axis,  $\gamma$  is inverted to the semicircle  $\overline{\gamma}$  erected on the  $x$ -axis and  $\mathcal{C}$  is inverted to the circle  $\overline{\mathcal{C}}$  tangent to  $\overline{\gamma}$  externally. We write the  $x$ -coordinates of the intersections of  $\overline{\alpha_i}, \overline{\alpha_j}$  and  $\overline{\gamma}$  with the  $x$ -axis as  $s, t$  and  $p, q$  with  $q < p$ . Then  $t < s$  since  $a_i < a_j$ .

By the definition of inversion we have

$$s = \frac{k^2}{2a_i}, t = \frac{k^2}{2a_j}, p = \frac{k^2}{2a}, q = -\frac{k^2}{2b}. \quad (1)$$




Figure 2

Since the  $x$ -coordinates of the center and the radius of  $\bar{C}$  are  $\frac{s+t}{2}$  and  $\frac{s-t}{2}$ , and those of  $\bar{\gamma}$  are  $\frac{p+q}{2}$  and  $\frac{p-q}{2}$ , we have

$$\left(\frac{s+t}{2} - \frac{p+q}{2}\right)^2 + d^2 = \left(\frac{s-t}{2} + \frac{p-q}{2}\right)^2,$$

where  $d$  is the  $y$ -coordinate of the center of  $\bar{C}$ . From this,

$$st - sp - tq + pq + d^2 = 0. \quad (2)$$

Since  $O$  is outside  $\bar{C}$ , we have

$$r = \frac{k^2}{\left| \left(\frac{s+t}{2}\right)^2 + d^2 - \left(\frac{s-t}{2}\right)^2 \right|} \cdot \frac{s-t}{2} = \frac{k^2}{\left(\frac{s+t}{2}\right)^2 + d^2 - \left(\frac{s-t}{2}\right)^2} \cdot \frac{s-t}{2}.$$

By using (1) and (2) we get the conclusion.  $\square$

**Lemma 2.** *If  $\alpha_i$  (resp.  $\alpha_j$ ) is the line, then the radius of the inscribed circle is*

$$\frac{-ab}{a_j - a} \text{ (resp. } \frac{ab}{a_i + b}).$$

*Proof.* Even in this case (2) in the proof of Lemma 1 holds with  $s = 0$  (resp.  $t = 0$ ), and we get the conclusion.  $\square$

**Theorem 3.** *Assume  $a \neq b$ , and let  $\alpha_i, \alpha_j \in \Phi(\alpha, \beta, \gamma)$  with  $\alpha_i < \alpha_j$ . The radius of the circle inscribed in the curvilinear triangle bounded by  $\alpha_i, \alpha_j$  and  $\gamma$  is*

$$\frac{ab(\mu(\alpha_i) - \mu(\alpha_j))}{b\mu(\alpha_i) - a\mu(\alpha_j)}.$$

*Proof.* If  $\alpha_i$  and  $\alpha_j$  are semicircles, then

$$\frac{ab(\mu(\alpha_i) - \mu(\alpha_j))}{b\mu(\alpha_i) - a\mu(\alpha_j)} = \frac{ab \left( \frac{a_i - a + b}{a_i} - \frac{a_j - a + b}{a_j} \right)}{b \cdot \frac{a_i - a + b}{a_i} - a \cdot \frac{a_j - a + b}{a_j}} = \frac{ab(a_j - a_i)}{a_i a_j - a a_i + b a_j}.$$

Hence the theorem follows from Lemma 1. If one of  $\alpha_i, \alpha_j$  is the line, the result follows from Lemma 2.  $\square$

Similarly we have

**Theorem 4.** Assume  $a = b$ , and let  $\alpha_i, \alpha_j \in \Phi(\alpha, \beta, \gamma)$  with  $\alpha_i < \alpha_j$ . The radius of the circle inscribed in the curvilinear triangle bounded by  $\alpha_i, \alpha_j$  and  $\gamma$  is

$$\frac{a^2(\mu(\alpha_j) - \mu(\alpha_i))}{a(\mu(\alpha_j) - \mu(\alpha_i)) - 1}.$$

The functions  $x \mapsto \frac{ab(1-x)}{b-ax}$ ,  $a \neq b$  and  $x \mapsto \frac{a^2x}{ax-1}$ ,  $a > 0$  are injective.

Therefore, we have

**Corollary 5.** Let  $\alpha_0, \alpha_1, \dots, \alpha_n \in \Phi(\alpha, \beta, \gamma)$  with  $\alpha_0 < \alpha_1 < \dots < \alpha_n$ . The circles inscribed in the curvilinear triangle bounded by  $\alpha_{i-1}, \alpha_i$  and  $\gamma$  ( $i = 1, 2, \dots, n$ ) are all congruent if and only if  $\mu(\alpha_0), \mu(\alpha_1), \dots, \mu(\alpha_n)$  is a geometric sequence if  $a \neq b$ , or an arithmetic sequence if  $a = b$ .

**Theorem 6.** Let  $\{\alpha_0 = \alpha, \alpha_1, \dots, \alpha_n = \beta, \gamma\}$  be an arbelos in  $n$ -aliquot parts. The common radius of the Archimedean circles in  $n$ -aliquot parts is

$$\begin{cases} \frac{ab \left( b^{\frac{2}{n}} - a^{\frac{2}{n}} \right)}{b^{\frac{2}{n}+1} - a^{\frac{2}{n}+1}}, & \text{if } a \neq b, \\ \frac{2a}{n+2}, & \text{if } a = b. \end{cases}$$

*Proof.* First we consider the case  $a \neq b$ . We can assume  $\alpha_0 < \alpha_1 < \dots < \alpha_n$  by renaming if necessary. The sequence  $\frac{b}{a} = \mu(\alpha_0), \mu(\alpha_1), \dots, \mu(\alpha_n) = \frac{a}{b}$  is a geometric sequence by Corollary 5. If we write its common ratio as  $d$ , we have  $\frac{a}{b} = d^n \left( \frac{b}{a} \right)$ , and then  $d = \left( \frac{a}{b} \right)^{\frac{2}{n}}$ . By Theorem 3 the radius of the Archimedean circle is

$$\frac{ab(1-d)}{b-ad} = \frac{ab \left( 1 - \left( \frac{a}{b} \right)^{\frac{2}{n}} \right)}{b - a \left( \frac{a}{b} \right)^{\frac{2}{n}}} = \frac{ab \left( b^{\frac{2}{n}} - a^{\frac{2}{n}} \right)}{b^{\frac{2}{n}+1} - a^{\frac{2}{n}+1}}.$$

Similarly we can get the second assertion.  $\square$

Note that the second assertion is the limiting case of the first assertion when  $b \rightarrow a$ .

**Theorem 7.** Let  $\{\alpha_0 = \alpha, \alpha_1, \dots, \alpha_n = \beta, \gamma\}$  be an arbelos in  $n$ -aliquot parts with  $\alpha_0 < \alpha_1 < \dots < \alpha_n$ . Then  $\alpha_i$  is the line in  $\Phi(\alpha, \beta, \gamma)$  if  $n$  is even and  $i = \frac{n}{2}$ .

Otherwise it is a semicircle with radius

$$\begin{cases} \left| \frac{b^{\frac{2i}{n}-1}(a-b)}{a^{\frac{2i}{n}-1} - b^{\frac{2i}{n}-1}} \right|, & \text{if } a \neq b, \\ \left| \frac{na}{n-2i} \right|, & \text{if } a = b. \end{cases}$$

*Proof.* Suppose  $a \neq b$ . Since  $\frac{b}{a} = \mu(\alpha_0)$ ,  $\mu(\alpha_1), \dots, \mu(\alpha_n) = \frac{a}{b}$  is a geometric sequence with common ratio  $\left(\frac{a}{b}\right)^{\frac{2}{n}}$ , we have  $\mu(\alpha_i) = \left(\frac{a}{b}\right)^{\frac{2i}{n}} \left(\frac{b}{a}\right) = \left(\frac{a}{b}\right)^{\frac{2i}{n}-1}$ .

If  $n$  is even and  $i = \frac{n}{2}$ , then  $\mu(\alpha_i) = 1$  and  $\alpha_i$  is the line. Otherwise,  $\mu(\alpha_i) \neq 1$  and  $\alpha_i$  is a semicircle. Let  $a_i$  be the  $x$ -coordinate of its center. The radius of  $\alpha_i$  is  $|a_i|$  and  $\frac{a_i - a + b}{a_i} = \left(\frac{a}{b}\right)^{\frac{2i}{n}-1}$ . From this,  $a_i = \frac{b^{\frac{2i}{n}-1}(a-b)}{b^{\frac{2i}{n}-1} - a^{\frac{2i}{n}-1}}$ .

The proof for the case  $a = b$  is similar.  $\square$

### 3. Embedded patterns of the arbelos

Let  $\{\alpha_0 = \alpha, \alpha_1, \dots, \alpha_n = \beta, \gamma\}$  be an arbelos in  $n$ -aliquot parts with  $\alpha_0 < \alpha_1 < \dots < \alpha_n$ . There exists a semicircle  $\gamma'$  which is tangent to all Archimedean circles externally. It is clearly concentric to  $\gamma$ . (If  $n = 1$  we will take for  $\gamma'$  the semicircle concentric to  $\gamma$  and tangent to the Archimedean circle externally). Let  $\alpha', \beta'$  be two semicircles in  $y \geq 0$ , tangent to  $\alpha_i$ s at  $O$  and also tangent to  $\gamma'$ . We take  $\alpha'$  in the region  $x \geq 0$  and  $\beta'$  in the region  $x \leq 0$ . Let  $a'$  and  $b'$  be the radii of  $\alpha'$  and  $\beta'$  respectively. Clearly  $\alpha', \beta'$  are tangent externally at  $O$ , and  $\gamma'$  intersects the  $x$ -axis at  $-2b'$  and  $2a'$ , and  $\Phi(\alpha, \beta, \gamma) \subseteq \Phi(\alpha', \beta', \gamma')$ . Moreover, for any  $\alpha_i \in \Phi(\alpha, \beta, \gamma)$ ,  $\mu(\alpha_i)$  considered in  $\Phi(\alpha, \beta, \gamma)$  is equal to  $\mu(\alpha_i)$  considered in  $\Phi(\alpha', \beta', \gamma')$  since the centers of  $\gamma$  and  $\gamma'$  coincide.

**Lemma 8.** (a) If  $a \neq b$ ,  $\left(\frac{a'}{b'}\right)^n = \left(\frac{a}{b}\right)^{n+2}$ .

(b) If  $a = b$ ,  $\frac{a'}{n} = \frac{a}{n+2}$ .

*Proof.* If  $a \neq b$  we have

$$\begin{aligned} a' &= a - \frac{ab \left( a^{\frac{2}{n}} - b^{\frac{2}{n}} \right)}{a^{\frac{2}{n}+1} - b^{\frac{2}{n}+1}} = \frac{a^{\frac{2}{n}+1} (a-b)}{a^{\frac{2}{n}+1} - b^{\frac{2}{n}+1}}, \\ b' &= b - \frac{ab \left( a^{\frac{2}{n}} - b^{\frac{2}{n}} \right)}{a^{\frac{2}{n}+1} - b^{\frac{2}{n}+1}} = \frac{b^{\frac{2}{n}+1} (a-b)}{a^{\frac{2}{n}+1} - b^{\frac{2}{n}+1}}, \end{aligned}$$

by the definitions of  $a'$  and  $b'$ . Then the the first assertion follows. The second assertion follows similarly.  $\square$

**Theorem 9.**  $\{\alpha', \alpha_0, \alpha_1, \dots, \alpha_n, \beta', \gamma'\}$  is an arbelos in  $(n+2)$ -aliquot parts.

*Proof.* Let us assume  $a \neq b$ . By Lemma 8 and the proof of Theorem 6,  $\mu(\alpha_0)$ ,  $\mu(\alpha_1), \dots, \mu(\alpha_n)$  is a geometric sequence with common ratio  $\left(\frac{a'}{b'}\right)^{\frac{2}{n+2}}$ . Also by Lemma 8 we have

$$\frac{\mu(\alpha_0)}{\mu(\alpha')} = \frac{b a'}{a b'} = \left(\frac{b'}{a'}\right)^{\frac{n}{n+2}} \frac{a'}{b'} = \left(\frac{a'}{b'}\right)^{\frac{2}{n+2}},$$

and

$$\frac{\mu(\beta')}{\mu(\alpha_n)} = \frac{a' b}{b' a} = \frac{a'}{b'} \left(\frac{b'}{a'}\right)^{\frac{n}{n+2}} = \left(\frac{a'}{b'}\right)^{\frac{2}{n+2}}.$$

The case  $a = b$  follows similarly.  $\square$

Let  $\{\alpha, \beta, \gamma\}$  be an arbelos and all the semicircles be constructed in  $y \geq 0$  such that the diameters lie on the  $x$ -axis. Let  $\alpha_{-1} = \alpha$ ,  $\alpha_1 = \beta$  and  $\gamma_1 = \gamma$ . If there exists an arbelos in  $(2n - 1)$ -aliquot parts  $\{\alpha_{-n}, \alpha_{-(n-1)}, \dots, \alpha_{-1}, \alpha_1, \dots, \alpha_n, \gamma_{2n-1}\}$  with  $\alpha_{-n} < \alpha_{-(n-1)} < \dots < \alpha_{-1} < \alpha_1 < \dots < \alpha_n$ , we shall construct an arbelos in  $(2n + 1)$ -aliquot parts as follows.

Let  $\gamma_{2n+1}$  be the semicircle concentric to  $\gamma$  and tangent externally to all Archimedean circles of the above arbelos. This meets the  $x$ -axis at two points one of which is in the region  $x > 0$  and the other in  $x < 0$ . We write the semicircle passing through  $O$  and the former point as  $\alpha_{-(n+1)}$  and the semicircle passing through  $O$  and the latter point as  $\alpha_{n+1}$ . Then  $\{\alpha_{-(n+1)}, \alpha_{-n}, \dots, \alpha_{-1}, \alpha_1, \dots, \alpha_{n+1}, \gamma_{2n+1}\}$  is an arbelos in  $(2n + 1)$ -aliquot parts by Theorem 9. Now we get the set of semicircles

$$\{\dots, \alpha_{-(n+1)}, \alpha_{-n}, \dots, \alpha_{-1}, \alpha_1, \dots, \alpha_n, \alpha_{n+1}, \dots, \gamma_1, \gamma_3, \dots, \gamma_{2n-1}, \dots\},$$

where  $\{\alpha_{-n}, \dots, \alpha_{-1}, \alpha_1, \dots, \alpha_n, \gamma_{2n-1}\}$  form the arbelos in  $(2n - 1)$ -aliquot parts for any positive integer  $n$ . We shall call the above configuration the *odd pattern*.

**Theorem 10.** *Let  $\delta_{2n-1}$  be one of the Archimedean circles in*

$$\{\alpha_{-n}, \alpha_{-(n-1)}, \dots, \alpha_{-1}, \alpha_1, \dots, \alpha_n, \gamma_{2n-1}\}.$$

*Then the radii of  $\alpha_{-n}$  and  $\alpha_n$  are*

$$\frac{a^{2n-1}(a-b)}{a^{2n-1}-b^{2n-1}} \quad \text{and} \quad \frac{b^{2n-1}(a-b)}{a^{2n-1}-b^{2n-1}},$$

*and the radii of  $\gamma_{2n-1}$  and  $\delta_{2n-1}$  are respectively*

$$\frac{(a^{2n-1} + b^{2n-1})(a-b)}{a^{2n-1} - b^{2n-1}} \quad \text{and} \quad \frac{a^{2n-1}b^{2n-1}(a-b)(a^2 - b^2)}{(a^{2n-1} - b^{2n-1})(a^{2n+1} - b^{2n+1})}.$$

*Proof.* Let  $\overline{a_{-n}}$  and  $\overline{a_n}$  be the radii of  $\alpha_{-n}$  and  $\alpha_n$  respectively. By Lemma 8 we have

$$\left(\frac{\overline{a_{-n}}}{\overline{a_n}}\right)^{\frac{1}{2n-1}} = \left(\frac{\overline{a_{-(n-1)}}}{\overline{a_{n-1}}}\right)^{\frac{1}{2n-3}} = \dots = \frac{\overline{a_{-1}}}{\overline{a_1}} = \frac{a}{b}. \quad (3)$$

Since  $\gamma_{2n-1}$  and  $\gamma$  are concentric, we have

$$\overline{a_{-n}} - \overline{a_n} = a - b. \quad (4)$$

By (3) and (4) we have

$$\begin{aligned} \overline{a_{-n}} &= \frac{a^{2n-1}(a-b)}{a^{2n-1} - b^{2n-1}}, \\ \overline{a_n} &= \frac{b^{2n-1}(a-b)}{a^{2n-1} - b^{2n-1}}. \end{aligned}$$

It follows that the radius of  $\gamma_{2n-1}$  is

$$\overline{a_{-n}} + \overline{a_n} = \frac{(a^{2n-1} + b^{2n-1})(a-b)}{a^{2n-1} - b^{2n-1}},$$

and that of  $\delta_{2n-1}$  is

$$\begin{aligned} &\frac{(a^{2n-1} + b^{2n-1})(a-b)}{a^{2n-1} - b^{2n-1}} - \frac{(a^{2n+1} + b^{2n+1})(a-b)}{a^{2n+1} - b^{2n+1}} \\ &= \frac{a^{2n-1}b^{2n-1}(a-b)(a^2 - b^2)}{(a^{2n-1} - b^{2n-1})(a^{2n+1} - b^{2n+1})}. \end{aligned}$$

□

As in the odd case, we can construct the *even* pattern of arbelos

$\{\dots, \beta_{-(n+1)}, \beta_{-n}, \dots, \beta_{-1}, \beta_0, \beta_1, \dots, \beta_n, \beta_{n+1}, \dots, \gamma_2, \gamma_4, \dots, \gamma_{2n}, \dots\}$   
inductively by starting with an arbelos in 2-aliquot parts  $\{\beta_{-1}, \beta_0, \beta_1, \gamma_2\}$ , where  $\beta_{-1} = \alpha$ ,  $\beta_1 = \beta$  and  $\gamma_2 = \gamma$ . By Theorem 9,  $\{\beta_{-n}, \dots, \beta_{-1}, \beta_0, \beta_1, \dots, \beta_n, \gamma_{2n}\}$  forms an arbelos in  $2n$ -aliquot parts for any positive integer  $n$ , and  $\beta_0$  is the line by Theorem 7. Analogous to Theorem 10 we have

**Theorem 11.** *Let  $\delta_{2n}$  be one of the Archimedean circles in*

$$\{\beta_{-n}, \beta_{-(n-1)}, \dots, \beta_{-1}, \beta_0, \beta_1, \dots, \beta_n, \gamma_{2n}\}.$$

*The radii of  $\beta_{-n}$  and  $\beta_n$  are*

$$\frac{a^n(a-b)}{a^n - b^n} \quad \text{and} \quad \frac{b^n(a-b)}{a^n - b^n},$$

*and the radii of  $\gamma_{2n}$  and  $\delta_{2n}$  are respectively*

$$\frac{(a^n + b^n)(a-b)}{a^n - b^n} \quad \text{and} \quad \frac{a^n b^n (a-b)^2}{(a^n - b^n)(a^{n+1} - b^{n+1})}.$$

**Corollary 12.** *Let  $c_n$  and  $d_n$  be the radii of  $\gamma_n$  and  $\delta_n$  respectively.*

$$\begin{aligned} a_n &= b_{2n-1}, \\ a_{-n} &= b_{-(2n-1)}, \\ c_{2n-1} &= c_{2(2n-1)}, \\ d_{2n-1} &= d_{4n-2} + d_{4n}. \end{aligned}$$

Figure 3 shows the even pattern together with the odd pattern reflected in the  $x$ -axis. The trivial case of these patterns can be found in [2].




Figure 3

#### 4. Some Applications

We give two applications here, with the same notations as in §3.

**Theorem 13.** *The external common tangent of  $\beta_n$  and  $\beta_{-n}$  touches  $\gamma_{4n}$  for any positive integer  $n$ .*

*Proof.* The distance between the external common tangents of  $\beta_n$  and  $\beta_{-n}$  and the center of  $\gamma_{2n}$  is  $\frac{\overline{b_n}^2 + \overline{b_{-n}}^2}{\overline{b_n} + \overline{b_{-n}}}$  where  $\overline{b_n}$  and  $\overline{b_{-n}}$  are the radii of  $\beta_n$  and  $\beta_{-n}$ . By

Theorem 11 this is equal to  $\frac{(a-b)(a^{2n} + b^{2n})}{a^{2n} - b^{2n}}$ , the radius of  $\gamma_{4n}$ .  $\square$

**Theorem 14.** *Let  $BK_n$  be the circle orthogonal to  $\alpha$ ,  $\beta$  and  $\delta_{2n-1}$ , and let  $AR_n$  be the inscribed circle of the curvilinear triangle bounded by  $\beta_n$ ,  $\beta_0$  and  $\gamma_{2n}$ . The circles  $BK_n$  and  $AR_n$  are congruent for every natural number  $n$ .*

*Proof.* Assume  $a \neq b$ . Since  $AR_n$  is the Archimedean circle of the arbelos in 2-aliquot parts  $\{\beta_{-n}, \beta_0, \beta_n, \gamma_{2n}\}$ , the radius of  $AR_n$  is

$$\frac{\overline{b_n} \overline{b_{-n}} (\overline{b_n} - \overline{b_{-n}})}{\overline{b_n}^2 - \overline{b_{-n}}^2} = \frac{a^n b^n (a - b)}{a^{2n} - b^{2n}},$$



by Theorem 6 and Theorem 11.

On the other hand  $BK_n$  is the inscribed circle of the triangle bounded by the three centers of  $\alpha$ ,  $\beta$ ,  $\delta_{2n-1}$ . Since the length of three sides of the triangle are  $a + d_{2n-1}$ ,  $b + d_{2n-1}$ ,  $a + b$ , the radius of  $BK_n$  is

$$\sqrt{\frac{abd_{2n-1}}{a+b+d_{2n-1}}} = \frac{a^n b^n (a-b)}{a^{2n} - b^{2n}},$$

by Theorem 10. □

This theorem is a generalization of Bankoff circle [1]. Bankoff's third circle corresponds to the case  $n = 1$  in this theorem.

## References

- [1] L. Bankoff, Are the twin circles of Archimedes really twins?, *Math. Magazine*, 47 (1974) 214–218.
- [2] H. Okumura, Circles patterns arising from results in Japanese geometry, *Symmetry: Culture and Science*, 8 (1997) 4–23.

Hiroshi Okumura: Department of Information Engineering, Maebashi Institute of Technology,  
460-1 Kamisadori Maebashi Gunma 371-0816, Japan  
*E-mail address:* okumura@maebashi-it.ac.jp

Masayuki Watanabe: Department of Information Engineering, Maebashi Institute of Technology,  
460-1 Kamisadori Maebashi Gunma 371-0816, Japan  
*E-mail address:* watanabe@maebashi-it.ac.jp