

## Applying Poncelet's Theorem to the Pentagon and the Pentagonal Star

Arthur Holshouser, Stanislav Molchanov, and Harold Reiter

**Abstract.** A special case of Poncelet's Theorem states that if all points on circle  $C_2$  lie inside of circle  $C_1$  and if a convex  $n$ -polygon,  $n \geq 3$ , or an  $n$ -star,  $n \geq 5$ , is inscribed in circle  $C_1$  and circumscribed about circle  $C_2$ , then there exists a family of such  $n$ -polygons and  $n$ -stars. Suppose all points on  $C_2$  lie inside of  $C_1$ ,  $R, r$ , are the radii of  $C_1, C_2$  respectively and  $\rho$  is the distance between the centers of  $C_1, C_2$ . For  $n \geq 3$ , in a companion paper we give an algorithm that computes the necessary and sufficient conditions on  $R, r, \rho$ , where  $R > r + \rho, r > 0$ , so that if we start at any arbitrary point  $Q$  on  $C_1$  and draw successive tangents to  $C_2$  (counterclockwise about the center of  $C_2$ ) then we will return to  $Q$  in exactly  $n$  steps and not return to  $Q$  in fewer than  $n$  steps. This will create the above family of  $n$ -polygons and  $n$ -stars. However, when  $n \geq 5$ , this companion paper relies on computers to find these conditions. In some ways, this is a sign of defeat. In this paper, we illustrate for  $n = 5$  a technique that can compute these exact same necessary and sufficient conditions on  $R, r, \rho$  without using a computer.

### 1. Introduction

A special case of Poncelet's Theorem states that if all points on circle  $C_2$  lie inside of circle  $C_1$  and if a convex  $n$ -polygon,  $n \geq 3$ , or an  $n$ -star,  $n \geq 5$ , is inscribed in circle  $C_1$  and circumscribed about circle  $C_2$  then there exists a family of such  $n$ -polygons and  $n$ -stars. Suppose all points on  $C_2$  lie inside of  $C_1$ ,  $R, r$  are the radii of  $C_1, C_2$  respectively and  $\rho$  is the distance between the centers of  $C_1, C_2$ .

For  $n = 5$ , we illustrate a technique that can be carried out by hand that computes the necessary and sufficient conditions on  $R, r, \rho$ , where  $R > r + \rho, r > 0$ , so that if we start at any point  $Q$  on  $C_1$ , and draw successive tangents to  $C_2$  (counterclockwise about the center of  $C_2$ ) then we will return to  $Q$  in exactly 5 steps and not return to  $Q$  in fewer than 5 steps.

If we consider  $R > \rho \geq 0$  to be arbitrary but fixed and consider  $r > 0$  to be a variable, then we end up with two polynomial equations  $P(R, \rho, r) = 0, \bar{P}(R, \rho, r) = 0$  that are each of third degree in the variable  $r$ . Each of the equations  $P(R, \rho, r) = 0, \bar{P}(R, \rho, r) = 0$  has exactly one  $r$ -root that satisfies  $R > r + \rho, r > 0$ . This  $r$ -root of  $P(R, \rho, r) = 0$  is the value of  $r$  so that we get

a family of pentagonal stars and this  $r$ -root of  $\bar{P}(R, \rho, r) = 0$  is the value of  $r$  so that we get a family of pentagons when we start at any arbitrary point  $Q$  on  $C_1$ .

In this paper, we only deal with the 5-star. The geometric reasoning for the convex pentagon is very similar. Also, we know from the companion paper [4] that the two polynomials  $P(R, \rho, r)$ ,  $\bar{P}(R, \rho, r)$  are related by

$$\bar{P}(R, \rho, r) = P(-R, \rho, r) = P(R, \rho, -r).$$

Thus, we can immediately write the polynomial  $\bar{P}(R, \rho, r)$  directly from the polynomial  $P(R, \rho, r)$  without doing any additional work.

**2. A preliminary unfactored form of the polynomial  $P(R, \rho, r)$**

In this section, for the pentagonal star, we compute a preliminary first version called  $P^*(R, \rho, r)$  of the polynomial  $P(R, \rho, r)$ . Then in Section 3, we refine  $P^*(R, \rho, r)$  by factoring it into four irreducible factors

$$P^*(R, \rho, r) = P(R, \rho, r) (2Rr + \theta) (r - R + \rho)^2,$$

where

$$P(R, \rho, r) = 8\rho^2 Rr^3 - 4R^2\theta r^2 - 2R\theta^2 r + \theta^3, \tag{1}$$

$$\theta = R^2 - \rho^2. \tag{2}$$

Of course, for the pentagon we have

$$\bar{P}(R, \rho, r) = P(-R, \rho, r) = P(R, \rho, -r) = -8\rho^2 Rr^3 - 4R^2\theta r^2 + 2R\theta^2 r + \theta^3.$$

The linear factor  $r - R + \rho = 0$  in  $P^*(R, \rho, 0)$  is extraneous since we require  $R > r + \rho, r > 0$ . Also, the factor  $2Rr + \theta = 0$  in  $P^*(R, \rho, r)$  is an Euler type of equation which has only an extraneous negative  $r$ -root since  $\theta > 0$ .




Figure 1. A drawing to compute  $P^*(R, \rho, r) = 0$

By Poncelet's Theorem, we can use any drawing to compute  $P^*(R, \rho, r) = 0$  that simplifies the problem. Therefore, by Poncelet's Theorem, the simple drawing of Figure 1 is all that we need to compute  $P^*(R, \rho, r) = 0$  for the pentagonal star. An analogous drawing is used for the convex pentagon. The  $\theta$  in Figure 1 is different from the  $\theta$  in (2)

$O_1$  is the center of the big circle  $C_1$  and  $O_2$  is the center of the inside circle  $C_2$ .  $R$  and  $r$  are the radii of  $C_1$  and  $C_2$  respectively, and  $\rho = O_1O_2$  is the distance between the centers  $O_1$  and  $O_2$ . We immediately have

$$(a) O_2y = (O_2v) \cdot \sin \frac{\theta}{2} = (R - \rho) \sin \frac{\theta}{2} = r,$$

$$(b) O_1t = (O_1w) \cdot \cos \phi = R \cos \phi = \rho + r.$$

The parametric equation of the line  $US$  is

$$\begin{cases} x = R \cos \phi - t(R \cos \phi + R \cos \theta), \\ y = -R \sin \phi + t(R \sin \phi + R \sin \theta), \quad t \in R. \end{cases}$$

From these,

$$\begin{aligned} & x(\sin \phi + \sin \theta) + y(\cos \phi + \cos \theta) \\ &= R \cos \phi(\sin \phi + \sin \theta) - R \sin \phi(\cos \phi + \cos \theta) = R \sin \theta \cos \phi - R \cos \theta \sin \phi \\ &= R \sin(\theta - \phi) \\ &= 2R \sin\left(\frac{\theta - \phi}{2}\right) \cos\left(\frac{\theta - \phi}{2}\right) \end{aligned}$$

Therefore,

$$\begin{aligned} & 2x \sin\left(\frac{\theta + \phi}{2}\right) \cos\left(\frac{\theta - \phi}{2}\right) + 2y \cos\left(\frac{\theta + \phi}{2}\right) \cos\left(\frac{\theta - \phi}{2}\right) \\ &= 2R \sin\left(\frac{\theta - \phi}{2}\right) \cos\left(\frac{\theta - \phi}{2}\right), \end{aligned}$$

and

$$x \sin \frac{\theta + \phi}{2} + y \cos \frac{\theta + \phi}{2} = R \sin \frac{\theta - \phi}{2}$$

is the equation of the line  $US$ .

Letting  $y = 0$  in this equation of the line  $US$ , we have

$$O_1x = \frac{R \sin \frac{\theta - \phi}{2}}{\sin \frac{\theta + \phi}{2}}.$$

Therefore,

$$xO_2 = O_1O_2 - O_1x = \rho - O_1x = \rho - \frac{R \sin \frac{\theta - \phi}{2}}{\sin \frac{\theta + \phi}{2}}.$$

Also,

$$O_2z = xO_2 \cdot \sin \frac{\theta + \phi}{2} = r = \left[ \rho - \frac{R \sin \frac{\theta - \phi}{2}}{\sin \frac{\theta + \phi}{2}} \right] \sin \frac{\theta + \phi}{2}.$$

Therefore,

$$\rho \sin \frac{\theta + \phi}{2} - R \sin \frac{\theta - \phi}{2} = r. \quad (3)$$

Now  $\sin^2 \frac{\theta + \phi}{2} = \frac{1 - \cos(\theta + \phi)}{2}$  and  $\sin^2 \frac{\theta - \phi}{2} = \frac{1 - \cos(\theta - \phi)}{2}$ . Also,  $\sin \frac{\theta + \phi}{2} \sin \frac{\theta - \phi}{2} = \frac{1}{2} \cos \phi - \frac{1}{2} \cos \theta$ . Squaring (3) and making these substitutions we have

$$\begin{aligned} \rho^2 [1 - \cos(\theta + \phi)] + R^2 [1 - \cos(\theta - \phi)] - 2\rho R [\cos \phi - \cos \theta] &= 2r^2, \\ -R^2 \cos(\theta - \phi) - \rho^2 \cos(\theta + \phi) + 2\rho R [\cos \theta - \cos \phi] &= 2r^2 - R^2 - \rho^2, \\ -R^2 [\cos \theta \cos \phi + \sin \theta \sin \phi] - \rho^2 [\cos \theta \cos \phi - \sin \theta \sin \phi] + 2\rho R [\cos \theta - \cos \phi] \\ &= 2r^2 - R^2 - \rho^2, \end{aligned}$$

Therefore,

$$(-R^2 + \rho^2) \sin \theta \sin \phi = 2r^2 - R^2 - \rho^2 - 2\rho R (\cos \theta - \cos \phi) + (R^2 + \rho^2) (\cos \theta \cos \phi).$$

Squaring we have

$$\begin{aligned} &(-R^2 + \rho^2)^2 (1 - \cos^2 \theta) (1 - \cos^2 \phi) \\ &= (-R^2 + \rho^2)^2 (1 - \cos \theta) (1 + \cos \theta) (1 - \cos \phi) (1 + \cos \phi) \quad (4) \\ &= [2r^2 - R^2 - \rho^2 - 2\rho R (\cos \theta - \cos \phi) + (R^2 + \rho^2) (\cos \theta \cos \phi)]^2. \end{aligned}$$

Since we have a homogeneous geometric equation in the variables  $R, r, \rho$ , it is convenient to let  $R = 1$ .

From (a), (b) we know that  $\sin \frac{\theta}{2} = \frac{r}{R - \rho} = \frac{r}{1 - \rho}$  and  $\cos \phi = \frac{\rho + r}{R} = \rho + r$ .

Therefore,  $\cos \theta = 1 - 2 \sin^2 \frac{\theta}{2} = 1 - 2 \left( \frac{r}{1 - \rho} \right)^2$  and  $\cos \phi = \rho + r$ . From these,

$$\begin{aligned} 1 - \cos \theta &= 2 \left( \frac{r}{1 - \rho} \right)^2, \\ 1 + \cos \theta &= 2 - 2 \left( \frac{r}{1 - \rho} \right)^2 = \frac{2(1 - \rho)^2 - 2r^2}{(1 - \rho)^2}, \\ 1 - \cos \phi &= 1 - \rho - r, \quad 1 + \cos \phi = 1 + \rho + r, \\ \cos \theta - \cos \phi &= 1 - \rho - r - 2 \left( \frac{r}{1 - \rho} \right)^2, \\ \cos \theta \cos \phi &= (\rho + r) \left( 1 - 2 \left( \frac{r}{1 - \rho} \right)^2 \right). \end{aligned}$$

If we make these substitutions and also put  $R = 1$  in (4), multiply the equation by  $(1 - \rho)^4$ , and partially simplify by straightforward calculations, transposing everything to one side of the equation, then we have the following equation which

we call the *preliminary polynomial equation*.

$$\begin{aligned}
& P^*(R, r, \rho) \\
&= \left[ (2r^2 + 2\rho r + \rho^2 - 2\rho - 1)(1 - \rho)^2 + 4\rho r^2 + (1 + \rho^2) \left( (1 - \rho)^2 - 2r^2 \right) (\rho + r) \right]^2 \\
&\quad - 4(1 - \rho^2)^2 \left[ r^2(1 - \rho - r)^2(1 - \rho + r)(1 + \rho + r) \right] \\
&= 0.
\end{aligned}$$

### 3. Factoring the preliminary equation $P^*(R, r, \rho) = 0$ into irreducible factors

The above preliminary polynomial equation  $P^*(R, r, \rho) = P^*(1, r, \rho) = 0$  in the variable  $r$  at first glance appears to be intractable. However, if we substitute specific values of  $\rho$ , e.g.  $\rho = 0, 1, 2$ , we quickly conjecture that this polynomial equation can probably be factored into simple factors.

If we substitute  $\rho = 0$ , the preliminary equation becomes

$$P^*(R, \rho, r) = P^*(1, 0, r) = [2r^2 - 1 + (1 - 2r^2)r]^2 - 4r^2(1 - r^2)^2 = 0,$$

which is equivalent to

$$\begin{aligned}
0 &= (2r^3 - 2r^2 - r + 1)^2 - (2r^3 - 2r)^2 \\
&= (-2r^2 + r + 1)(4r^3 - 2r^2 - 3r + 1) \\
&= -(r - 1)(2r + 1)(r - 1)(4r^2 + 2r - 1) \\
&= -(r - 1)^2(2r + 1)(4r^2 + 2r - 1).
\end{aligned}$$

By making other substitutions for  $\rho$ , we soon conjecture that

$$P^*(R, r, \rho) = P^*(1, r, \rho) = P(1, r, \rho)(2r + 1 - \rho^2)(r - 1 + \rho)^2 = 0$$

where  $P(1, r, \rho)$  is a 3rd degree polynomial in  $r$ .

We now rigorously prove this conjecture. By direct substitution of  $r = 1 - \rho$  into  $P^*(1, r, \rho)$  we can easily prove that  $r = 1 - \rho$  is a double  $r$ -root of  $P^*(1, r, \rho) = 0$ . To see this, we see that  $P^*(1, r, \rho)$  is of the form

$$P^* = [\text{xxx}]^2 - [\text{yyy}](1 - \rho - r)^2$$

and we only need to show that  $[\text{xxx}] = 0$  when  $r = 1 - \rho$  to show that  $r = 1 - \rho$  is a double root of  $P^*(1, r, \rho) = 0$ .

Now in  $[\text{xxx}]$  when  $r = 1 - \rho$  we see that

$$\begin{aligned}
2r^2 + 2\rho r + \rho^2 - 2\rho - 1 &= 2r(\rho + r) + \rho^2 - 2\rho - 1 \\
&= 2(1 - \rho) + \rho^2 - 2\rho - 1 \\
&= \rho^2 - 4\rho + 1.
\end{aligned}$$

Therefore, in [xxx] when  $r = 1 - \rho$  we have

$$\begin{aligned} & (2r^2 + 2\rho r + \rho^2 - 2\rho - 1)(1 - \rho)^2 + 4\rho r^2 \\ &= (\rho^2 - 4\rho + 1)(1 - \rho)^2 + 4\rho(1 - \rho)^2 \\ &= (\rho^2 + 1)(1 - \rho)^2. \end{aligned}$$

Also, in [xxx] when  $r = 1 - \rho$ , we have

$$\begin{aligned} (1 + \rho^2)((1 - \rho)^2 - 2r^2)(\rho + r) &= (1 + \rho^2)((1 - \rho)^2 - 2(1 - \rho)^2) \\ &= -(1 + \rho^2)(1 - \rho)^2. \end{aligned}$$

Therefore, when  $r = 1 - \rho$ ,

$$[\text{xxx}] = (\rho^2 + 1)(1 - \rho)^2 - (1 + \rho^2)(1 - \rho)^2 = 0,$$

and  $r = 1 - \rho$  is a double  $r$ -root of  $P^*(1, r, \rho) = 0$ .

The proof that  $2r + 1 - \rho^2 = 0$  gives an  $r$ -root of  $P^*(1, r, \rho) = 0$  takes a little longer but it is completely straightforward.

Therefore, we know that

$$P^*(1, r, \rho) = (ar^3 + br^2 + cr + d)(2r + 1 - \rho^2)(r - 1 + \rho)^2$$

where  $a, b, c, d$  need to be determined.

Rewriting  $P^*(1, r, \rho) = a_0r^6 + a_1r^5 + a_2r^4 + a_3r^3 + a_4r^2 + a_5r + a_6$ , it is fairly easy by straightforward calculations to compute the following coefficients.

$$\begin{aligned} a_0 &= 16\rho^2, \\ a_1 &= -8(1 - \rho)(-\rho^3 + 3\rho^2 + \rho + 1), \\ a_5 &= -2(1 - \rho)^5(1 + \rho)^4, \\ a_6 &= (1 - \rho)^6(1 + \rho)^4. \end{aligned}$$

As an example, we have  $a_0 = 4(1 + \rho^2)^2 - 4(1 - \rho^2)^2 = 16\rho^2$ . Also, to compute  $a_5$  we have the following relevant terms,

$$\begin{aligned} & \left( 2\rho(1 - \rho)^2 r + (\rho^2 - 2\rho - 1)(1 - \rho)^2 + (1 + \rho^2)(1 - \rho)^2 r + (1 + \rho^2)(1 - \rho)^2 \rho \right)^2 \\ &= (2\rho(1 - \rho)^2 r + (1 + \rho^2)(1 - \rho)^2 r + (\rho^2 - 2\rho - 1)(1 - \rho)^2 + (1 + \rho^2)(1 - \rho)^2 \rho)^2 \\ &= ((1 + \rho)^2(1 - \rho)^2 r + (\rho^3 + \rho^2 - \rho - 1)(1 - \rho)^2)^2 \\ &= ((1 + \rho)^2(1 - \rho)^2 r - (1 + \rho)^2(1 - \rho)^3)^2. \end{aligned}$$

From this, we see that  $a_5 = -2(1 - \rho)^5(1 + \rho)^4$ .

To compute  $a_6$  we let  $r = 0$  in  $P^*(1, r, \rho)$  and we have

$$\begin{aligned} a_6 &= ((\rho^2 - 2\rho - 1)(1 - \rho)^2 + (1 + \rho^2)(1 - \rho)^2\rho)^2 \\ &= (1 - \rho)^4(\rho^3 + \rho^2 - \rho - 1)^2 \\ &= (1 - \rho)^4((\rho + 1)^2(\rho - 1))^2 \\ &= (1 - \rho)^6(1 + \rho)^4. \end{aligned}$$

The calculation of  $a_1$  is a little longer but it is completely straightforward. However, we must be careful not to overlook anything in computing  $a_1$ . Once we know  $a_0, a_1, a_5, a_6$ , it is completely straight forward to compute

$$P(1, r, \rho) = ar^3 + br^2 + cr + d = 8\rho^2r^3 - 4\theta r^2 - 2\theta^2r + \theta^3,$$

where  $\theta = 1 - \rho^2$ . So  $P^*(1, r, \rho) = P(1, r, \rho) \cdot (2r + 1 - \rho^2)(r - 1 + \rho)^2$ . We now proceed to rigorously prove this. We first note that

$$(8\rho^2r^3 - 4\theta r^2 - 2\theta^2r + \theta^3)(2r + \theta) = 16\rho^2r^4 - 8\theta^2r^3 - 8\theta^2r^2 + \theta^4.$$

Therefore, we prove that

$$\begin{aligned} P^*(1, r, \rho) &= (16\rho^2r^4 - 8\theta^2r^3 - 8\theta^2r^2 + \theta^4)(r - 1 + \rho)^2 \\ &= (16\rho^2r^4 - 8\theta^2r^3 - 8\theta^2r^2 + \theta^4)(r^2 - 2(1 - \rho)r + (1 - \rho)^2). \end{aligned}$$

This equality will be true if and only if the equality correctly computes the above values for  $a_0, a_1, a_5, a_6$ , since we have already proved that  $2r + \theta$  and  $(r - 1 + \rho)^2$  are factors of  $P^*(1, r, \rho)$ . Now  $a_0 = 16\rho^2$  is obviously computed correctly. Also,

$$\begin{aligned} a_1 &= -32\rho^2(1 - \rho) - 8(1 - \rho^2)^2 \\ &= -8(1 - \rho)(4\rho^2 + (1 + \rho)^2(1 - \rho)) \\ &= -8(1 - \rho)(-\rho^3 + 3\rho^2 + \rho + 1), \\ a_5 &= -2(1 - \rho)(1 - \rho^2)^4 \\ &= -2(1 - \rho)^5(1 + \rho)^4, \\ a_6 &= (1 - \rho^2)^4(1 - \rho)^2 \\ &= (1 - \rho)^6(1 + \rho)^4. \end{aligned}$$

Therefore, we have now rigorously proved that

$$\begin{aligned} P^*(1, r, \rho) &= P(1, r, \rho)(2r + \theta)(r - 1 + \rho)^2 \\ &= (8\rho^2r^3 - 4\theta r^2 - 2\theta^2r + \theta^3)(2r + \theta)(r - 1 + \rho)^2 \end{aligned}$$

where  $\theta = 1 - \rho^2$ .

Of course, this equation can be written for  $P^*(R, r, \rho)$  in the three variables  $R, r, \rho$  since the equation is a homogeneous geometric equation. This equation  $P(R, r, \rho) = P(1, r, \rho)$  is exactly the same equation that we derived in a companion paper by using a computer. This computer derivation was carried out independently by Prof. Benjamin Klein of Davidson College and by Parker Garrison. So we now have three independent verifications of this one equation.

#### 4. Studying $P(R, r, \rho) = P(1, r, \rho)$

If  $R = 1 > \rho \geq 0$ , we require  $R = 1 > r + \rho, r > 0$ .

It is easy to show that  $P(1, r, \rho) = 8\rho^2 r^3 - 4\theta r^2 - 2\theta^2 r + \theta^3$  is irreducible in the rational field.

Letting  $R = 1, 0 < \rho < 1$ , we know by Descartes's law of signs that  $P(1, r, \rho) = 0$  has two or zero positive  $r$ -roots for each fixed  $0 < \rho < 1$ . For each fixed  $0 < \rho < 1$  we show that  $P(1, r, \rho) = 0$  has one  $r$ -root that satisfies  $0 < r < 1 - \rho$ . ( $\rho = 0$  is easy to deal with.)

Now  $P(1, r, \rho) = P(1, +\infty, \rho) > 0$ .

Also,  $P(1, r, \rho) = P(1, 0, \rho) > 0$ . If we show that  $P(1, r, \rho) = P(1, 1 - \rho, \rho) < 0$ , then it will follow that for each fixed  $0 < \rho < 1$ ,  $P(1, r, \rho) = 0$  will have one  $r$ -root that satisfies  $0 < r < 1 - \rho$ .

Now  $P(1, r, \rho) = P(1, 1 - \rho, \rho) < 0$  if and only if

$$(1 - \rho)^3(8\rho^2 - 4(1 + \rho) - 2(1 + \rho)^2 + (1 + \rho)^3) < 0.$$

This is true since

$$\begin{aligned} & (1 - \rho)^3(-4(1 + \rho - 2\rho^2) - (1 + \rho)^2(2 - (1 + \rho))) \\ &= (1 - \rho)^3(-4(1 + 2\rho)(1 - \rho) - (1 + \rho)^2(1 - \rho)) \\ &= -(1 - \rho)^4(4(1 + 2\rho) + (1 + \rho)^2) \\ &< 0. \end{aligned}$$

Therefore, for each  $R = 1 > \rho \geq 0$ , we see that  $P(1, r, \rho) = 0$  has one  $r$ -root that satisfies  $R = 1 > r + \rho, r > 0$ .

If  $R = 1 > \rho \geq 0$  are fixed, this  $r$ -root is the radius of the inside circle  $C_2$  so that we have a family of 5-stars that are inscribed in  $C_1$  and circumscribed about  $C_2$  when the distance between the centers of  $C_1, C_2$  is  $\rho$ ,  $R = 1$  is the radius of  $C_1$  and  $r$  is the radius of  $C_2$ .

#### 5. Extending the equation to include convex pentagons

By using analogous reasoning we can show that the companion equation

$$\bar{P}(R, r, \rho) = P(-R, r, \rho) = P(R, -r, \rho) = -8\rho^2 R r^3 - 4R^2 \theta r^2 + 2R\theta^2 r + \theta^3 = 0,$$

where  $\theta = R^2 - \rho^2$  is the relation between  $R, r, \rho, R > r + \rho, r > 0$ , so that we have a family of convex pentagons that are inscribed in  $C_1$  and circumscribed about  $C_2$ . From the companion paper, we know that the equation  $\bar{P}(R, r, \rho) = 0$  for the convex pentagon can be written directly from  $\bar{P}(R, r, \rho) = P(-R, r, \rho) = P(R, -r, \rho)$ . It is easy to show that for  $R = 1 > \rho \geq 0$ , there exists exactly one real  $r$ -root of  $\bar{P}(1, r, \rho) = 0$  that satisfies  $R = 1 > r + \rho, r > 0$ .

#### 6. Concluding remarks

Everything in this paper was done completely by hand and this adds completeness to a computer only derived solution. The advantage of the computer derived solution is that it is less mentally demanding and requires less thought to carry out.



## References

- [1] N. A. Court, *College Geometry*, Dover reprint, 2007.
- [2] V. Dragović and M. Radnović, *Poncelet Porisms and Beyond, Integrable Billiards, Hyperelliptic Jacobians and Pencils of Quadrics*, Springer, 2011;  
[http://www.springer.com/cda/content/document/cda\\_downloaddocument/9783034800143-p1.pdf?SGWID=0-0-45-1054140-p174034876](http://www.springer.com/cda/content/document/cda_downloaddocument/9783034800143-p1.pdf?SGWID=0-0-45-1054140-p174034876).
- [3] L. Halbeisen, <https://www.math.uzh.ch/fileadmin/math/.../01.14.p...>, University of Zurich.
- [4] A. Holshouser, S. Molchanov, and H. Reiter, A special case of Poncelet's problem, *Forum Geom.*, 16 (2016) 151–170.
- [5] E. Weisstein, Poncelet's Porism,  
<http://mathworld.wolfram.com/PonceletsPorism.html>.

Arthur Holshouser: 3600 Bullard St., Charlotte, North Carolina, USA

*E-mail address:* xxx

Stanislav Molchanov: Department of Mathematics, University of North Carolina Charlotte, Charlotte, North Carolina 28223, USA

*E-mail address:* smolchan@uncc.edu

Harold Reiter: Department of Mathematics, University of North Carolina Charlotte, Charlotte, North Carolina 28223, USA

*E-mail address:* hbreiter@email.uncc.edu